

HILGARDIA

A JOURNAL OF AGRICULTURAL SCIENCE

PUBLISHED BY THE

CALIFORNIA AGRICULTURAL EXPERIMENT STATION

VOL. 2

FEBRUARY, 1927

No. 10

EFFECT OF MULCHES ON SOIL TEMPERATURES DURING THE WARMEST WEEK IN JULY, 1925

ALFRED SMITH*

Previous investigations^{1, 2, 3} on the use of paper as a mulch in contrast to soil mulches have dealt mainly with the effect on crop growth and crop yields. Investigations are now under way regarding the effects of such paper covering on the temperature and moisture conditions of the soil, and also on the most desirable kind of paper to use with particular regard to color, weight, durability, and need for perforations.

By the use of an unperforated black paper mulch at Berkeley in 1924, Shaw⁴ found that the soil at a depth of three inches averaged about 0.42° F warmer than at a similar depth in the soil-mulched plots. Hartung² in the pineapple fields of Hawaii obtained higher mean soil temperatures at a depth of three inches in areas covered with grayish brown paper mulches than in unprotected soil. In his summary he states that "paper mulch maintains a mean soil temperature in the upper 3-inch layer of soil during the cool season in the localities given, from 3 to 4.5 degrees Fahrenheit above that of non-paper covered soil; provided the mulch paper is dark, preferably black in color." Although the standard mulch paper which Hartung used presented a "greyish brown appearance," he nevertheless recommends black paper. Stewart⁵ and his co-workers in Hawaii found that on clear days the areas covered with black paper were from 12° to 15° F warmer during the day, and from 4° to 5° F warmer during the night.

* Assistant Professor of Soil Technology and Associate Soil Technologist in the Experiment Station.



Fig. 1.—Soil temperature plots at Davis, California, January, 1925.

METHODS

The experimental tract used in these experiments is an area of Yolo loam which had been summer fallowed in 1924. The surface soil to a depth of three feet is a fairly uniform loam on a subsoil of fine sandy loam with some minor variations of coarser texture. This soil is of recent alluvial deposition, is derived mainly from sedimentary rocks, and occupies a nearly flat topographic position. Sixteen plots, each five meters square, were arranged in this area, separated by paths two meters wide.

A sixteen-point recorder was installed in a one-story frame structure, 150 feet north of the experimental area, in January, 1925, with all connecting wires carried to the plots in overhead conduits as shown




Fig. 2.—Automatic temperature recorder.

in figure 1. Sixteen resistance thermometers were carefully standardized against standard mercury thermometers and were placed in the soil at the center of only five of the plots at depths of from $\frac{1}{2}$ inch to 36 inches. The least number of thermometers in any one plot was two and the most was six. In burying the thermometers, a hole of small diameter was dug, the soil of the various horizons being carefully laid aside. The thermometers were then inserted in the undisturbed soil on the north side of the holes, so that the resistance units or "bulbs" were at least eight inches from the excavation. The hole was then filled, the soil layers being put back in proper order and lightly tamped in order to attain the same degree of compactness as existed originally.

The temperature recorder (fig. 2), a Leeds & Northrup 16-point recorder, was adjusted so that the temperature of each individual thermometer was recorded every 15 minutes, giving 96 records in a 24-hour period or 10,752 records from the 16 thermometers in a period of 7 days. All temperatures herein reported are in degrees Fahrenheit.

A continuous record of the air temperature and humidity was obtained in a standard United States Weather Bureau shelter, located in the northeast corner of the area.

SURFACE TREATMENTS OF CERTAIN PLOTS IN 1925

Various treatments of the soil surface were under experiment in 1925, but only plots 6, 7, 10, 11 and 15, in which soil temperatures were taken, will be discussed in this paper. The surface treatments of these plots were as follows:

Plot 6—covered with Pabco Thermo-Gen 214, black on both sides and with large, irregularly triangular perforations $1\frac{1}{4}$ inches apart.

Plot 7—not covered with paper, cultivated 4 inches deep once a month.

Plot 10—covered with Moistite Thermo-Gen mulch paper, gray on both sides and with small triangular perforations $\frac{3}{4}$ of an inch apart.

Plot 11—covered with Thermo-Gen, black on both sides, but without perforations.

Plot 15—covered with Mulch Paper Plain, gray on both sides and without perforations.

Where the paper mulch was put on, it was laid in 36-inch strips running north and south. A lap of 3 inches was allowed and over this redwood battens were laid and stapled down with No. 6 iron wire.

These battens were $1\frac{1}{4}$ inches wide and $\frac{1}{4}$ inch thick (fig. 3). All plots were covered on May 1 and during the 1925 season no crop was grown on any of them.

A period of very high temperatures with clear weather occurred in California, centering around July 16-18, 1925. At some points in the state all previous maximum temperatures for July were equalled or exceeded. The discussion in this paper is confined to a consideration of the temperature changes recorded during this week, or from July 14 to July 21.



Fig. 3.—Paper mulch and bare plots, Davis, California, May, 1925.

EXPERIMENTAL RESULTS

In two plots (Nos. 6 and 7) temperatures were obtained at the depth of one-half inch. The average day temperature (sunrise to sunset) in the bare plot, No. 7, at this depth was 109° , and in plot 6, which was covered with black perforated paper, it was 99° , or 10 degrees colder. The night (sunset to sunrise) temperatures were reversed, however, the average of the bare plot being 81° and of the covered plot 86.6° , or a difference in favor of the covered plot of 5.6° . The highest soil temperature recorded in any of these plots in 1925 was 143° obtained at a depth of one-half inch in the bare plot (No. 7) on July 17.

At the six-inch depth in these same two plots the average day temperature was 91.3° for the bare plot and 90.4° for the covered plot, or a difference of only 0.9° in favor of the bare plot. At the

same depth the average night temperature was slightly higher in both cases, it being 92.7° for the bare plot and 92.1° for the covered plot.

A more detailed study of the records from the bare plot at the various depths shows that the average day and night temperatures (fig. 4) to a depth of twelve inches during this warm period differ decidedly; but at greater depths little or no difference was noted. This is in full agreement with the records for any of the clear days in 1925 not herewith reported, as temperatures were obtained from February to October.




Fig. 4.—Temperature ranges in bare plot, July 14-21, 1925.

The average time of sunrise during the week of July 14-21 was 4:54, and for sunset it was 7:30. The time of occurrence of the maximum and minimum temperatures in the bare plot are shown in table 1. There was no diurnal effect below 12 inches, but there was a gradual increase in temperature, which at the 24-inch depth amounted during the week to four degrees. The maximum air temperature for the week occurred at approximately the same time as that at the one-half inch depth, while the minimum air temperature occurred 28 minutes earlier than the minimum at the one-half inch depth.

The rate at which the heat moved into the soil is shown graphically in figure 5 and the loss of heat from the soil is illustrated in figure 6.

TABLE 1
TIME OF OCCURRENCE OF MAXIMUMS AND MINIMUMS IN BARE PLOT

Depth	Time of maximum after sunrise	Time of minimum after sunset
½ inch.....	8 hours 51 minutes.....	8 hours 51 minutes
3 inches.....	11 hours.....	10 hours 43 minutes
6 inches.....	12 hours 17 minutes.....	11 hours 51 minutes
12 inches.....	16 hours 25 minutes.....	14 hours

The gradual increase in temperatures at depths below 12 inches shows that heat was not only lost by radiation from the soil to the atmosphere, but also by conduction downward.

The diurnal range in temperature in the bare plot for depths to 12 inches naturally varies inversely to the depth and is shown in figure 7.

When the temperatures at the 3 inch depth in all five plots are compared (figs. 8, 9), it is quite evident that different results were obtained.




Fig. 5.—Maximum temperatures in bare plot: time of occurrence after sunrise. Average for the week of July 14-21, 1925.




Fig. 6.—Minimum temperatures in bare plot: time of occurrence after sunset. Average for the week of July 14-21, 1925.




Fig. 7.—Average diurnal range for the week of July 14-21, 1925.




Fig. 8.—Temperatures at 3 inch depth in bare and covered plots and of air by two hour intervals. Week of July 14-21, 1925.




Fig. 9.—Temperatures at 3 inch depth in bare and covered plots and of air by two hour intervals. Week of July 14-21, 1925.

TABLE 2
AVERAGE TEMPERATURES AT THREE-INCH DEPTH, JULY 14-21, 1925

Plot	Cover	Day	Night	Maximum	Minimum
		° F.	° F.	° F.	° F.
11	Black solid paper.....	98.0	96.0	107.0	88.7
7	No paper—soil mulch.....	92.6	89.6	100.5	83.4
6	Black perforated paper.....	91.4	90.4	97.9	84.7
10	Gray perforated paper.....	86.3	85.0	92.4	79.4
15	Gray solid paper.....	88.0	86.0	94.0	81.3
Air			102.0	66.9

In table 2 the effect of the color of the paper mulch as well as the effect of perforations is shown. Plot 11, which was covered with black solid paper, was 5.4° warmer during the day and 6.4° warmer during the night than the bare plot (No. 7). Where the black perforated paper was used as a mulch (plot 6), the average day and night temperatures were about the same as in the bare plot, with slightly higher night temperatures in favor of the black perforated paper.

Plot 15, which was covered with gray solid paper, was 4.6° colder during the day and 3.6° colder during the night than the bare plot. The coldest of the five plots was number 10, where the average day temperature was 6.3° lower than in the bare plot (No. 7). As the color of the surface soil in these plots is brown, drying to a light brown, the plots covered with the light-colored or gray papers did not have the capacity to absorb as much heat as the bare plot, while those covered with black paper had an increased capacity for heat absorption.

The effect of perforating the mulch paper with large and frequent perforations (fig. 10) is to permit the circulation of air, which results in general in lower temperatures being recorded at the 3 inch depth in the plots covered with perforated paper than in those covered with the non-perforated paper. Plot 6, which was covered with black perforated paper, was on the average 1.2° colder during the day and slightly warmer during the night, as shown in table 2, than the bare plot. The gray perforated paper (plot 10) was the coldest of the five plots, both during the day and at night. Whatever heat is absorbed by the paper mulches during periods of sunlight is rapidly lost from the soil surface by conduction and radiation to the atmosphere when the papers are perforated.

The maximum temperatures in the five plots at the 3-inch depth occurred usually at about the same time, which was two hours after the maximum air temperature was reached. The average maximums




Fig. 10.—Types of perforated paper used.

of these plots as given in table 2 show the range of from 94° to 107° , or 13° . The minimum temperatures at the same depth occurred on the average 1 hour and 40 minutes after the minimum air temperature and showed a range from 81.3° to 88.7° , or 7.4° .

The changes in temperature which occurred at the 12 inch depth in these five plots are graphically shown in figures 11 and 12. The maximum temperatures occurred usually at about the same time, which was 8 hours after the maximum air temperature was reached, and showed a range of 11.6° . The minimums occurred, on the average, 6 hours after the minimum air temperatures and had a range of 10.6° . The results for the 12 inch depth are shown in table 3.

TABLE 3
AVERAGE TEMPERATURES AT TWELVE-INCH DEPTH, JULY 14-21, 1925

Plot	Day	Night	Maximum	Minimum
	$^{\circ} F.$	$^{\circ} F.$	$^{\circ} F.$	$^{\circ} F.$
11.....	93.1	95.4	96.0	92.6
7.....	87.6	89.4	90.3	86.4
6.....	88.6	90.0	90.6	87.7
10.....	82.7	84.0	84.4	82.0
15.....	84.0	85.0	85.6	83.0
Air.....	102.0	66.9




Fig. 11.—Temperatures at 12 inch depth in bare and covered plots and of air by two hour intervals. Week of July 14-21, 1925.




Fig. 12.—Temperatures at 12 inch depth in bare and covered plots and of air by two hour intervals. Week of July 14-21, 1925.

In general the effects of the various types of mulches were the same at a depth of 12 inches (table 3) as those at the 3 inch depth. Plot 11, covered with black solid paper, had the highest temperatures at the 12 inch depth. The average day temperature was 5.5° , and the average night temperature 6.0° , higher than in the bare plot. Plot 6, covered with black perforated paper, was slightly warmer at the 12 inch depth than the bare plot. The plots covered with gray paper mulches were the coldest at the 12 inch depth. Plot 15 covered with gray solid paper was 3.6° and 4.4° colder on the average during the day and night, respectively, than the bare plot (No. 7). Plot 10, covered with the gray perforated paper, was on the average during the day 4.9° and during the night 5.4° colder than the bare plot.

SUMMARY

The highest soil temperature found was in the bare plot, where on July 17, 1925, at a depth of one-half inch, 143° F was registered. Temperatures obtained at a depth of one-half inch showed that the bare plot averaged 10° warmer during the day and 5.6° cooler at night than the plot covered with perforated black paper. In the bare plot the average day temperature at the 6 inch depth for the week was 0.9° higher and the average night temperature was 0.6° higher than on the area covered with perforated black mulch paper.

In the bare plot, where temperatures were obtained at depths of $\frac{1}{2}$, 3, 6, 12, 24, and 36 inches, decided differences were found between the night and day temperatures to a depth of 12 inches.

Temperatures taken at a depth of 3 inches in five plots, where different mulches were used, varied considerably. The warmest plot was that covered with solid black paper. Where the perforated black paper was used the temperatures were about the same as in the bare plot. The coldest plots were those covered with gray paper and in this case the perforated paper was again colder than the non-perforated. The maximum temperatures at the 3 inch depth occurred on the average 2 hours after the maximum air temperature and showed a range of 13° . The minimum temperatures at the 3 inch depth occurred on the average 1 hour and 40 minutes after the minimum air temperature and had a range of 7.4° .

Temperatures taken at a depth of 12 inches showed that the plot covered with the solid black paper was the warmest. The plot covered with perforated black paper was slightly warmer at the 12 inch depth

than the bare plot. The coldest plot was that covered with gray perforated paper, while the gray solid paper was only slightly warmer than the gray perforated. The maximum temperatures at the 12 inch depth occurred usually about 8 hours after the maximum air temperature was reached and showed a range in the various plots of 11.6°. The minimum temperatures at the same depth occurred, on the average, 6 hours after the minimum air temperature and had a range of 10.6°.

The effect of the mulch paper was markedly influenced by variations in color and by the presence or absence of perforations. The warmest soil during this week was that covered with non-perforated black paper, and the coldest that covered with the gray perforated paper. Under the conditions of this experiment, the standard perforated mulch papers showed no material effect in increasing the soil temperatures.

These results are not given as indicating the possible effects which might occur when paper mulches are used in crop rows and are not to be taken as recommendations for the use of any particular type of paper mulch. Other work is in progress which will include the effects on crops. This will be reported later.

LITERATURE CITED

¹ ECKART, CHAS. F.

1923. How thermogen enhances the growth of plants 1-29. Published by author, San Francisco.

² HARTUNG, W. J.

1926. The functions of paper mulch in pineapple culture 1-31. Hawaiian Pineapple Co., Ltd., Honolulu.

³ LYON, H. L., G. R. STEWART, and WALLACE MACFARLANE

1923. Univ. Hawaii, second annual short course in pineapple production: 36-39, 74-75, 143-144.

⁴ SHAW, C. F.

1926. The effect of a paper mulch on soil temperature. *Hilgardia* 1: 341-365.

⁵ STEWART, G. R., E. C. THOMAS, and JOHN HORNER

1926. Some effects of mulching paper on Hawaiian soils. *Soil Sci.* 22: 35-51.

Transmitted September, 1926.

PUBLICATIONS AVAILABLE FOR FREE DISTRIBUTION

BULLETINS

- | No. | No. |
|--|--|
| 253. Irrigation and Soil Conditions in the Sierra Nevada Foothills, California. | 870. Browning of Yellow Newtown Apples. |
| 261. Melaxuma of the Walnut, "Juglans regia." | 871. The Relative Cost of Yarding Small and Large Timber. |
| 262. Citrus Diseases of Florida and Cuba Compared with Those of California. | 872. The Cost of Producing Market Milk and Butterfat on 246 California Dairies. |
| 263. Size Grades for Ripe Olives. | 873. Pear Pollination. |
| 268. Growing and Grafting Olive Seedlings. | 874. A Survey of Orchard Practices in the Citrus Industry of Southern California. |
| 273. Preliminary Report on Kearney Vineyard Experimental Drain. | 875. Results of Rice Experiments at Cor-tena, 1923. |
| 275. The Cultivation of Belladonna in California. | 876. Sun-Drying and Dehydration of Wal-nuts. |
| 276. The Pomegranate. | 877. The Cold Storage of Pears. |
| 277. Sudan Grass. | 879. Walnut Culture in California. |
| 278. Grain Sorghums. | 880. Growth of Eucalyptus in California Plantations. |
| 279. Irrigation of Rice in California. | 881. Growing and Handling Asparagus Crowns. |
| 283. The Olive Insects of California. | 882. Pumping for Drainage in the San Joaquin Valley, California. |
| 294. Bean Culture in California. | 883. Monilia Blossom Blight (Brown Rot) of Apricot. |
| 304. A Study of the Effects of Freezes on Citrus in California. | 885. Pollination of the Sweet Cherry. |
| 310. Plum Pollination. | 886. Pruning Bearing Deciduous Fruit Trees. |
| 312. Mariout Barley. | 887. Fig Smut. |
| 313. Pruning Young Deciduous Fruit Trees. | 888. The Principles and Practice of Sun-drying Fruit. |
| 319. Caprifigs and Caprification. | 889. Berseem or Egyptian Clover. |
| 324. Storage of Perishable Fruit at Freez-ing Temperatures. | 890. Harvesting and Packing Grapes in California. |
| 325. Rice Irrigation Measurements and Experiments in Sacramento Valley, 1914-1919. | 891. Machines for Coating Seed Wheat with Copper Carbonate Dust. |
| 328. Prune Growing in California. | 892. Fruit Juice Concentrates. |
| 331. Phylloxera-Resistant Stocks. | 893. Crop Sequences at Davis. |
| 335. Coconut Meal as a Feed for Dairy Cows and Other Livestock. | 894. Cereal Hay Production in California. Feeding Trials with Cereal Hay. |
| 339. The Relative Cost of Making Logs from Small and Large Timber. | 895. Bark Diseases of Citrus Trees. |
| 340. Control of the Pocket Gopher in California. | 896. The Mat Bean (Phaseolus aconitifolius). |
| 343. Cheese Pests and Their Control. | 897. Manufacture of Roquefort Type Cheese from Goat's Milk. |
| 344. Cold Storage as an Aid to the Mar-keting of Plums. | 898. Orchard Heating in California. |
| 346. Almond Pollination. | 899. The Blackberry Mite, the Cause of Redberry Disease of the Himalaya Blackberry, and its Control. |
| 347. The Control of Red Spiders in Decid-uous Orchards. | 900. The Utilization of Surplus Plums. |
| 348. Pruning Young Olive Trees. | 901. Cost of Work Horses on California Farms. |
| 349. A Study of Sidedraft and Tractor Hitches. | 902. The Codling Moth in Walnuts. |
| 350. Agriculture in Cut-over Redwood Lands. | 903. Farm-Accounting Associations. |
| 352. Further Experiments in Plum Pollina-tion. | 904. The Dehydration of Prunes. |
| 353. Bovine Infectious Abortion. | 905. Citrus Culture in Central California. |
| 354. Results of Rice Experiments in 1922. | 906. Stationary Spray Plants in California. |
| 357. A Self-mixing Dusting Machine for Applying Dry Insecticides and Fungicides. | 907. Yield, Stand and Volume Tables for White Fir in the California Pine Region. |
| 358. Black Measles, Water Berries, and Related Vine Troubles. | 908. Alternaria Rot of Lemons. |
| 361. Preliminary Yield Tables for Second Growth Redwood. | 909. The Digestibility of Certain Fruit By-products as Determined for Rumi-nants. |
| 362. Dust and the Tractor Engine. | 910. Factors Affecting the Quality of Fresh Asparagus after it is Harvested. |
| 363. The Pruning of Citrus Trees in Cali-fornia. | 911. Paradichlorobenzene as a Soil Fumi-gant. |
| 364. Fungicidal Dusts for the Control of Bunt. | 912. A Study of the Relative Values of Cer-tain Root Crops and Salmon Oil as Sources of Vitamin A for Poultry. |
| 365. Avocado Culture in California. | 913. The California Poultry Industry; a Statistical Study. |
| 366. Turkish Tobacco Culture, Curing and Marketing. | 914. Planting and Thinning Distances for Deciduous Fruit Trees. |
| 367. Methods of Harvesting and Irrigation in Relation of Mouldy Walnuts. | |
| 368. Bacterial Decomposition of Olives dur-ing Pickling. | |
| 369. Comparison of Woods for Butter Boxes. | |

CIRCULARS

- | | |
|--|---|
| <p>No.</p> <p>87. Alfalfa.</p> <p>117. The Selection and Cost of a Small Pumping Plant.</p> <p>127. House Fumigation.</p> <p>129. The Control of Citrus Insects.</p> <p>136. <i>Melilotus indica</i> as a Green-Manure Crop for California.</p> <p>144. Oidium or Powdery Mildew of the Vine.</p> <p>157. Control of the Pear Scab.</p> <p>160. Lettuce Growing in California.</p> <p>164. Small Fruit Culture in California.</p> <p>166. The County Farm Bureau.</p> <p>170. Fertilizing California Soils for the 1918 Crop.</p> <p>173. The Construction of the Wood-Hoop Silo.</p> <p>178. The Packing of Apples in California.</p> <p>179. Factors of Importance in Producing Milk of Low Bacterial Count.</p> <p>190. Agriculture Clubs in California.</p> <p>199. Onion Growing in California.</p> <p>202. County Organizations for Rural Fire Control.</p> <p>203. Peat as a Manure Substitute.</p> <p>209. The Function of the Farm Bureau.</p> <p>210. Suggestions to the Settler in California.</p> <p>212. Salvaging Rain-Damaged Prunes.</p> <p>215. Feeding Dairy Cows in California.</p> <p>217. Methods for Marketing Vegetables in California.</p> <p>220. Unfermented Fruit Juices.</p> <p>228. Vineyard Irrigation in Arid Climates.</p> <p>230. Testing Milk, Cream, and Skim Milk for Butterfat.</p> <p>231. The Home Vineyard.</p> <p>232. Harvesting and Handling California Cherries for Eastern Shipment.</p> <p>234. Winter Injury to Young Walnut Trees during 1921-22.</p> <p>235. Soil Analysis and Soil and Plant Inter-relations.</p> <p>236. The Common Hawks and Owls of California from the Standpoint of the Rancher.</p> <p>237. Directions for the Tanning and Dressing of Furs.</p> <p>238. The Apricot in California.</p> <p>239. Harvesting and Handling Apricots and Plums for Eastern Shipment.</p> <p>240. Harvesting and Handling Pears for Eastern Shipment.</p> <p>241. Harvesting and Handling Peaches for Eastern Shipment.</p> <p>243. Marmalade Juice and Jelly Juice from Citrus Fruits.</p> <p>244. Central Wire Bracing for Fruit Trees.</p> <p>245. Vine Pruning Systems.</p> <p>247. Colonization and Rural Development.</p> <p>248. Some Common Errors in Vine Pruning and Their Remedies.</p> <p>249. Replacing Missing Vines.</p> <p>250. Measurement of Irrigation Water on the Farm.</p> <p>252. Supports for Vines.</p> <p>253. Vineyard Plans.</p> <p>254. The Use of Artificial Light to Increase Winter Egg Production.</p> | <p>No.</p> <p>255. Leguminous Plants as Organic Fertilizer in California Agriculture.</p> <p>256. The Control of Wild Morning Glory.</p> <p>257. The Small-Seeded Horse Bean.</p> <p>258. Thinning Deciduous Fruits.</p> <p>259. Pear By-products.</p> <p>261. Sewing Grain Sacks.</p> <p>262. Cabbage Growing in California.</p> <p>263. Tomato Production in California.</p> <p>264. Preliminary Essentials to Bovine Tuberculosis Control.</p> <p>265. Plant Disease and Pest Control.</p> <p>266. Analyzing the Citrus Orchard by Means of Simple Tree Records.</p> <p>267. The Tendency of Tractors to Rise in Front: Causes and Remedies.</p> <p>269. An Orchard Brush Burner.</p> <p>270. A Farm Septic Tank.</p> <p>272. California Farm Tenancy and Methods of Leasing.</p> <p>273. Saving the Gophered Citrus Tree.</p> <p>274. Fusarium Wilt of Tomato and its Control by Means of Resistant Varieties.</p> <p>276. Home Canning.</p> <p>277. Head, Cane, and Cordon Pruning of Vines.</p> <p>278. Olive Pickling in Mediterranean Countries.</p> <p>279. The Preparation and Refining of Olive Oil in Southern Europe.</p> <p>281. The Results of a Survey to Determine the Cost of Producing Beef in California.</p> <p>282. Prevention of Insect Attack on Stored Grain.</p> <p>283. Fertilizing Citrus Trees in California.</p> <p>284. The Almond in California.</p> <p>285. Sweet Potato Production in California.</p> <p>286. Milk Houses for California Dairies.</p> <p>287. Potato Production in California.</p> <p>288. Phyloxera Resistant Vineyards.</p> <p>289. Oak Fungus in Orchard Trees.</p> <p>290. The Tangier Pea.</p> <p>291. Blackhead and Other Causes of Loss of Turkeys in California.</p> <p>292. Alkali Soils.</p> <p>293. The Basis of Grape Standardization.</p> <p>294. Propagation of Deciduous Fruits.</p> <p>295. The Growing and Handling of Head Lettuce in California.</p> <p>296. Control of the California Ground Squirrel.</p> <p>298. The Possibilities and Limitations of Coöperative Marketing.</p> <p>299. Poultry Breeding Records.</p> <p>300. Coccidiosis of Chickens.</p> <p>301. Buckeye Poisoning of the Honey Bee.</p> <p>302. The Sugar Beet in California.</p> <p>303. A Promising Remedy for Black Measles of the Vine.</p> <p>304. Drainage on the Farm.</p> <p>305. Liming the Soil.</p> <p>306. A General Purpose Soil Auger and its Use on the Farm.</p> <p>307. American Foulbrood and its Control.</p> <p>308. Cantaloupe Production in California.</p> |
|--|---|

The publications listed above may be had by addressing

College of Agriculture,
University of California,
Berkeley, California.